

BalanceTutor™

Protocol & Clinical Guidelines

Oct 2016



BalanceTutor a dynamic and static postural control trainer

Contents

1 INTRODUCTION	3
2 NEUROMUSCULAR FUNCTION	4
3 NEUROMUSCULAR DYSFUNCTION	5
3.1 NEUROLOGICAL PROBLEMS	5
3.2 ORTHOPEDIC PROBLEMS	5
3.3 VESTIBULAR PROBLEMS	5
3.4 SPORT INJURIES	6
4 BALANCETUTOR SYSTEM	7
5 SAFETY	8
5.1 FORBIDDEN USE – CONTRAINDICATIONS	8
6 PROTOCOLS & APPLICATIONS	10
6.1 NEUROLOGICAL REHABILITATION	10
6.1.1 <i>Stroke</i>	10
6.1.2 <i>Multiple Sclerosis</i>	17
6.1.3 <i>Parkinson’s disease</i>	21
6.1.4 <i>Falls Prevention</i>	26
6.2 ORTHOPEDIC AND SPORT REHABILITATION	31
7 REFERENCE	49
8 VERSION HISTORY	56



1 Introduction

A postural perturbation is a sudden change in conditions that displaces the body posture away from equilibrium. Reactive postural control response is an automatic “like reflex” response which is initiated by unexpected postural perturbation in order to maintain balance and can be improved through practice.

Traditional physical rehabilitation is based mainly on proactive training (self-initiated actions). Clinical research has shown that in addition to proactive training, reactive training must be targeted. Traditional tools and methods simply can’t target reactive response training due to major safety issues and a lack of sophisticated treatment customization.

The system’s unique technology for the first time allows training reactive postural control response while standing or walking at different gait phases. Together with proactive training the system allows for optimal rehabilitation outcome and recovery.

2 Neuromuscular Function

Neuromuscular function relates to the ability of the body to efficiently recruit a muscle or group of muscles in order to perform a specific task and works on two main categories:

Interlimb coordination - Interlimb coordination primarily involves movements requiring sequential and simultaneous use of both sides of the body with a high degree of rhythmicity.

Intralimb coordination - Interlimb coordination is commonly divided into two categories: bimanual coordination and coordination of hands/feet.

For instance, coordination between adequate muscle activities of plantar flexors of one leg together with adequate magnitude of the ground reaction force provided by contralateral limb during heel strike is critical for normal gait ability and Balance.

In addition to Interlimb coordination (between limbs) proper Intralimb coordination (within limb) is also needed for normal gait ability and Balance. For instance, Spatio-temporal parameters or muscle synergies of several joints of the limb such as ankle, knee and hip joints is critical for Intralimb coordination during standing or walking. The Intralimb coordination can be impaired due to inadequate Spatio-temporal parameters leading to compensatory strategies adopted by post stroke patients.

Example I - Stepping forward requires coordination of weight bearing on one limb and Center of Mass transferring by another limb in initial swing phase in interlimb coordination.

Example II - Stepping forward requires coordination of Hip and knee flexion of the limb in initial swing phase.

3 Neuromuscular Dysfunction

Neuromuscular function can be impaired as a result of varied body system dysfunction as follows;

3.1 Neurological problems

Neurological disease and injuries such as Stroke, Head injuries, Spinal cord injury, Cerebral Palsy and Parkinson's disease can affect gait and balance ability. The features of Neurological problems vary according to medical condition, type of rehabilitation received, and other individual differences.

The gait of neurological patients is mainly characterized by reduced speed, stride length, and cadence, decreased angular excursions at leg joints, increased energetic cost, and asymmetry in kinematic and kinetic variables

These kinematic and kinetic deficits caused by neurological problems must be specifically targeted and properly rehabilitated for achieving optimal physical rehabilitation.

Normal balance ability and gait control requires adequate proactive and reactive postural control responses. Proactive response refers to ability of the body to maintain balance before some perturbation to the balance of the body such as a voluntary step whereas reactive postural response refers to ability of the body to maintain the balance of the body after unexpected perturbation like a compensatory step following a slip or trip.

Effective rehabilitation program must include both proactive and reactive postural control response evaluation and treatment.

Unfortunately, traditional rehabilitation is mostly based on proactive balance control response and not reactive responses. This limited rehabilitation program of traditional therapy mainly arises from safety issues and lack of treatment customization.

BalanceTutor system enables to target both proactive and reactive response to achieve optimal rehabilitation using professional protocols as demonstrated in this guide.

3.2 Orthopedic problems

Orthopedics disease and injuries such as Joint surgery, Amputation, Prosthetic, Muscle weakness, Ligament sprain and Muscle/ tendon strain can affect gait and balance ability. The features of orthopedic problems vary according to medical condition, type of rehabilitation received, and other individual differences.

3.3 Vestibular problems

The vestibular system is the sensory system that provides the leading contribution about the sense of balance and spatial orientation for the purpose of coordinating movement with balance. As movements consist of rotations and translations, the vestibular system comprises two components: the semicircular canal system, which indicates rotational movements; and the otoliths, which indicate linear accelerations. The vestibular system sends signals primarily to the neural structures

that control eye movements, and to the muscles to keep a body upright. The projections to the former provide the anatomical basis of the vestibulo-ocular reflex, which is required for clear vision; and the projections to the muscles that control posture are necessary to keep an animal upright.

The brain uses information from the vestibular system in the head and from proprioception throughout the body to understand the body's dynamics and kinematics (including its position and acceleration) from moment to moment.

Vestibular function together with proper function of proprioceptive sensors located in soft tissues such as muscles, tendon and ligament can be damaged due to varied of diseases or injuries and even by aging.

3.4 Sport injuries

Sports injuries are commonly caused by overuse, direct impact, or the application of force that is greater than the body part can structurally withstand. Some of the more common sports injuries include ankle sprain, Knee injuries, Hamstring and Groin strain.

Following a musculoskeletal damage due to sport injuries the proper function of proprioceptive sensors located in soft tissue can be affected leading to deficit of postural control and balance ability.

Proper balance ability requires adequate proactive and reactive postural control responses. Proactive response refers to ability of the body to prepare before an anticipates perturbation like hitting a ball whereas reactive response refers to ability of the body to regain balance after an unexpected perturbation like a compensatory step following a slip or trip.

In regards to motor learning, it is well known that by specific task practicing the reactive automatic response can be learned and improved. Therefore, exposing the athlete to an unexpected perturbation in safe environment is needed to train properly neuromuscular system for optimal postural control and balance ability achievement.

A dedicated proprioceptive rehabilitation program based on both proactive and reactive balance training is needed to regain optimal postural control and balance ability after the injury and reduce significantly recurrent sport injuries.



4 BalanceTutor System

The BalanceTutor by MediTouch is a Perturbation Treadmill that is an innovative system for postural control and balance training. The device consists of a treadmill mounted on a moving force plate platform. The platform moves in medial/ lateral and forward/ backward direction to simulate a slip and a trip in both the standing and walking phase.

An innovative approach using wireless sensors integrated in the BalanceTutor allows for the provision of numerous kinds of controlled expected and unexpected perturbation. These different kinds of perturbation can be generated in relation to the specific phase of the gait namely in stance or swing phase. In addition to controlled expected and unexpected perturbation in the standing and gait, the device also allows for Center of Pressure control practice.

5 Safety

In general, any kind of training on the system requires special attention to the ability and medical condition of the patient. However, the following list highlights particular indications that require specific care in use according to relevant therapy protocols and must be with the approval of a qualified physician:

- Joint instability following surgery or trauma
- Soft tissue damage such as Muscle or ligaments tear
- ACL/PCL restoration of Knee
- Droop foot
- Limb prosthesis

5.1 Forbidden use – Contraindications

Any kind of medical problems including cardiovascular, mental or physical impairment leading to an inability to use the system are contraindicated for the device. The following contraindications must be taken into account before using of the BalanceTutor:

- ! Patients who are not able to stand or walk without walk aid devices or external assistance.
- ! Body weight greater than 135 kg
- ! Severely fixed contractures
- ! Bone instability (non-consolidated fractures, unstable spinal column, severe osteoporosis)
- ! Open skin lesions in the area of the lower limbs and torso
- ! Unstable circulation
- ! Cardiac (blood) contraindications
- ! Uncooperative or (self) aggressive behavior, such as transitory psychotic syndrome
- ! Severe cognitive deficits
- ! Patients with (long-term) infusions
- ! Mechanical ventilation
- ! Severe vascular disorders of the lower limbs
- ! In general, patients who have been ordered to remain in bed or immobile
- ! Joint replacement like Hip, knee and Ankle.

The above list does not claim to be exhaustive. The decision as to whether a patient is suitable for treatment always comes under the remit of the physician in charge, who has sole medical



responsibility for the treatment. As part of this, he must evaluate in particular, in each individual case, possible risks and side-effects of the treatment against the benefit gained from it. In addition, the patient's individual situation plays just as important a role as the basic risk assessment for specific patient groups.

Being a scientific discipline, medicine is subject to constant change in response to new knowledge and progress. It is therefore the task of the physician in charge to continually keep his knowledge up to date by reading the latest scientific literature and to acquire new knowledge during the course of treatment.

6 Protocols & Applications

When an unexpected perturbation occurs, Center of Mass (COM) passively moves toward the perturbation's direction. The body needs to react immediately (70-120 ms) to restrain this unplanned COM passive movement. This reactive postural response is intended to stabilize the COM over the Base of Support by returning it back to the initial position or to create new safe BOS for COM using a compensatory step. In these two cases the body tries to move actively to the opposite direction of the applied perturbation.

There are several guidelines in this document that are intended, based on the concept above, to explain how to formulate the appropriate perturbation according to the rehabilitation aims as follows. Furthermore, perturbation can also be triggered at any phase of the gait cycle based on kinetics or kinematics parameters. Based on these guidelines the therapist can customize the treatment as demonstrated in the following protocols:

6.1 Neurological Rehabilitation

Numerous neurological diseases and injuries such as Stroke, Parkinson, Head injuries, Cerebral Palsy, Multiple Sclerosis and Spinal Cord Injuries can be treated by BalanceTutor as detailed in the following paragraphs.

6.1.1 Stroke

Falls are the number one medical complication after acute stroke [1–2]. Furthermore, the high fall risk for individuals with stroke is not only present in the acute phase, but it remains a considerable health concern throughout the poststroke life span. Because the incidence and prevalence of stroke increase as a result of ageing of the population [3] and the prevalence also increases as a result of continued improvement of post-stroke life expectancy [4], the societal impact of falls in stroke is rapidly growing. The impact is primarily related to the physical and psychosocial consequences of falls, which can be devastating. For instance, post stroke individuals are much more likely to sustain a hip fracture due to a fall and more often lose independent mobility and suffer from a higher mortality rate after a hip fracture [5–6]. These findings make falls and their prevention an important issue for every person involved in stroke care (neurologists, physiatrists, physiotherapists, nurses, and also caregivers at home) and all of the post stroke stages.

Both prospective and retrospective studies have consistently reported high fall rates in individuals with stroke. For comparison, in the general population of elderly people, ~30 percent fall at least once a year and ~15 percent fall twice or more [7–9], yielding an incidence rate of ~0.65 falls each person-year [10]. In the limited time that individuals with stroke are admitted in an acute care setting, 3.8 to 22.0 percent of the patients fall at least once [1–2,11], which makes falling the most frequent medical complication during hospitalization after stroke (~40% of all complications). The variability in these fall rates can be attributed at least partly to differences in length of stay. Expressed as fall incidence rates, they vary considerably less between studies, yielding 2.2 to 4.9 falls per person a year [1–2, 11]. Compared with other pathologies, such as congestive heart failure and community-acquired pneumonia, the risk for hospital falls is more than doubled in acute stroke [1]. The incidence rates are remarkably high given the degree of surveillance and the proportion of patients bedridden with, consequently, little exposure to risky situations.


The reported rates of people falling during inpatient rehabilitation range from 10.5 to 47.0 percent [12–24], with 5 to 27 percent of patients falling twice or more [14–16, 18–19]. Patients are most likely to fall during the first 3 weeks of rehabilitation [20–21]. The proportion of fallers as well as fall incidence rates (1.3–6.5 falls each person year) [14–16,18–21] varies considerably between studies, but particularly the incidence rates are, without exception, much higher than in the general population of elderly people.

In community-dwelling of stroke survivors, fall incidents are very common as well. The proportion of fallers increase as times goes by: 23-34, 40-73 and 43-70 percent for 3-4 month [12, 25], 6-month [17, 26–33], and 1-year follow-up [22, 34–36], respectively. Fallers in the stroke population are also more likely to become repeat fallers in comparison to elderly people in the general population. Most studies report proportions of repeat fallers in stroke populations between 21-57 percent for a 6-12 month period [17, 22, 27–28, 30–36]. Most studies have identified balance and gait deficits as important fall risk factors [17, 19, 27, 29, 31, 33, and 35].

Furthermore, research has suggested that individuals with stroke are more likely to fall when walking as it requires substantial cognitive control (i.e., was less automated), because fallers are more often unable to walk and talk at the same time or slow down when performing a concurrent mental task [28,35,44].

BalanceTutor system allows for several unique reactive response training in both standing and walking using controlled perturbation in simple task or multitask performance as follows:

6.1.1.1 Weight bearing on paretic limb – Medial / Lateral




Before perturbation




After perturbation

Patient position:	Standing comfortably with minimal external support
Platform's direction:	moves from healthy to paretic side
Comments:	partial weight bearing - low intensity and full weight bearing - high intensity

6.1.1.2 Weight bearing on paretic limb- Anterior/ Posterior




Before perturbation



After perturbation

Patient position:	Comfortable forward stance of paretic limb with minimal external support
Platform's direction:	Anterior to posterior
Comments:	partial weight bearing - low intensity and full weight bearing - high intensity

6.1.1.3 Equilibrium training




Before perturbation




After perturbation

Patient position:	Decreased base of support while standing with minimal external support
Platform's direction:	Anterior to posterior
Comments:	partial weight bearing - low intensity and full weight bearing - high intensity

6.1.1.4 Knee snapping training




Before perturbation



After perturbation

Patient position:	Standing at forward stance with weight bearing on slightly flexed paretic leg
Platform's direction:	posterior to Anterior
Comments:	partial weight bearing - low intensity and full weight bearing - high intensity

6.1.1.5 Compensatory step of paretic leg training



Before perturbation



After perturbation

Patient position:	Comfortable walking speed with minimal external support
Platform's direction:	Perturbation at swing phase of paretic leg
Comments:	For early stage of rehabilitation - Medial/ Lateral from healthy to paretic side and for advanced stage of rehabilitation - Medial/ Lateral from paretic to healthy side

6.1.1.6 Stabilization of paretic leg




Before perturbation




After perturbation

Patient position:	Comfortable walking speed with minimal external support
Platform's direction:	Medial/ Lateral Perturbation at stance phase of paretic leg
Comments:	For early stage of rehabilitation – Medial / Lateral from healthy to paretic side and for advanced stage of rehabilitation – Medial / Lateral from paretic to healthy side

6.1.1.7 Hip extensor muscle strengthening



Before perturbation




After perturbation

Patient position: Standing comfortably with minimal external support

Platform's direction: posterior to Anterior

Comments: This treatment can also be used while walking

6.1.1.8 Drop foot training



Before perturbation




After perturbation

Patient position: Standing comfortably with minimal external support


Platform's direction: posterior to Anterior

Comments: Fast and high level muscle contraction of ankle extensors

6.1.1.9 Unique upper paretic limb muscle activation




Before perturbation




After perturbation

Patient position:	Standing comfortably with minimal external support
Platform's direction:	Medial / Lateral from healthy to paretic side
Comments:	This technique 'forces' the patient to initiate an active movement of upper limb

6.1.1.10 Neuromuscular coordination



Before perturbation



After perturbation

Patient position:	Comfortable walking speed with minimal external support
Platform's direction:	Medial / Lateral Perturbation at stance and swing phase of paretic leg
Comments:	For early stage of rehabilitation – Medial / Lateral from healthy to paretic side and for advanced stage of rehabilitation – Medial / Lateral from paretic to healthy side

6.1.2 Multiple Sclerosis


Multiple sclerosis (MS) is a chronic progressive disease and the most common neurologic disease of young adults.

Existing research indicates that people with MS fall frequently [54–56], likely more than the general population [57–59], and more than the elderly, a group in which fall risk [60–16] and its effects are well studied and characterized [62–68]. A cross-sectional study in Italy found that 54 percent (27/50) of their small sample of subjects reported falling at least once and 32 percent (16/50) reported falling twice or more within the previous 2 months [54]. Similarly, another larger cross-sectional study of falls among people with MS aged 45 to 90 years in the United States (n = 1,089) found that 52 percent of subjects reported falling at least once in the previous 6 months [55]. Most recently, the first prospective study of falls among people with MS found that 63 percent of the subjects (48/76) recorded at least one fall and 58 percent (44/76) recorded two or more falls over a 3 month period [56]. Peterson et al. found that, of 354 people with MS aged 55 to 94 years in the United States, more than 50 percent reported receiving medical care for a fall-related injury at least once and 12 percent reported receiving medical care for a fall-related injury in the 6 months before the interview [69]. This potential for fall-related injuries likely contributes to loss of independence (mobility and activities of daily living) as well as reduced length [70] and quality of life [55] for people with MS.


Falls in people with MS have been found to be associated with being male [55–56], impaired balance [54–56], reduced ability to walk [54–56, 72], and use of a cane [54] or other walking aid [56]. In addition, studies have identified disturbed proprioception [56, 72], spasticity [55,72], and more severe MS (higher Expanded Disability Status Scale scores) [56] as risk factors for falls. People with MS also report that divided attention, reduced muscular endurance, fatigue, and heat sensitivity can cause them to fall [72]. Establishing whether individuals with MS have more injurious falls than other groups and identifying high-risk groups with MS is essential to developing effective and meaningful interventions.

BalanceTutor system allows for several unique postural control ability assessments as well as reactive response training in both standing and walking using controlled perturbation in simple task or multitask performance as follows:

6.1.2.1 Gait Ataxia training



Before perturbation




After perturbation

Patient position: Comfortable walking speed with minimal external support


Platform's direction: Anterior to posterior

Comments: Increasing walking speed according to patient ability

6.1.2.2 Dysmetria



Before perturbation




After perturbation

Patient position: Standing comfortably with minimal external support


Platform's direction: Medial / Lateral and Anterior/Posterior in four directions

Comments: Challenging and forcing the patient to react by initiating a compensatory step while walking

6.1.2.3 Hypertonia




Before perturbation



After perturbation

Patient position:	Standing comfortably with minimal external support
Platform's direction:	Medial/ Lateral and Anterior/Posterior in four directions
Clinical effects:	Mild perturbation in order to cause ankle and hip balance strategy and improvement of Automatic Postural Adjustment and controlled posture

6.1.2.4 Proprioceptive Rehabilitation




Before perturbation




After perturbation

Patient position:	Comfortable walking speed with minimal external support
Platform's direction:	Random utilizing four directions
Comments:	This treatment is suitable while standing and walking with customized mild perturbation

6.1.2.5 Vestibular Ataxia




Before perturbation



After perturbation

Patient position:	Comfortable walking speed with minimal external support
Platform's direction:	Random utilizing four directions
Comments:	This treatment is suitable while standing and walking with customized mild perturbation with rotation of the head during the treatment

6.1.2.6 Eye Movement Abnormalities




Before perturbation



After perturbation

Patient position:	Comfortable walking speed with minimal external support
Platform's direction:	Random utilizing four directions
Comments:	This treatment is suitable while standing and walking with customized mild perturbation with gaze fixation at different targets during the treatment

6.1.2.7 Neuromuscular coordination



Before perturbation



After perturbation

Patient position:	Comfortable walking speed with minimal external support
Platform's direction:	Random utilizing four directions
Clinical effects:	Multi-joint coordination and Coordinated Muscle group activation


6.1.3 Parkinson's disease

Parkinson's disease (PD) is a chronic and degenerative neurological disorder that leads to performance changes that affect an individual's ability to stay active PD results in severe disability in approximately 1 percent of people over the age of 60 [73–74], including veterans. The Department of Veterans Affairs treats an estimated 40,000 veterans with PD each year [75]. Akinesia, which typically occurs about 10 years after onset [76–77], is one of the most debilitating symptoms of PD. Studies report prevalence of akinesia ranging from 32 [76] to 60 percent [78] among people with PD. Akinesia is noted most prominently in the gait pattern of the person with PD. Akinetic gait is seen as a series of shuffling steps with reduced step length [79–80] and sometimes includes moments when movement ceases completely. These moments are referred to as “frozen gait,” which is generally described as feeling as if the feet are glued to the floor [67, 78, and 80]. Freezing is most common in initiating gait, turning, and walking through doorways or narrow spaces [74,76–78,81–82]. In addition to physical location, freezing is influenced by the person's emotions, appearing more frequently in times of distress [81] or when the person is attempting complex tasks, such as doing two things at once [82]. Freezing gait also increases the risk for falls [73– 74, 77, 81]. Gray and Hildebrand examined 118 people with PD and found that freezing is closely linked to reports of falling. Of patients with occasional or frequent episodes of freezing, 80 percent experienced falls”. Greater depression and lesser quality of life [83] are also found among those who have a history of falls, gait disturbances,

and akinesia and rigidity [84], especially if they subsequently experience further decline in physical mobility, emotional reactions, pain, and social isolation [85]. People with akinesia due to PD often continue to experience gait impairment even with current medical treatments [90,86]. As a result, nonpharmacological approaches are needed for akinesia, including the use of external sensory cues to help the person initiate or maintain movement. People with PD use a wide variety of techniques to overcome akinesia, including being pushed, marching to a cadence, rocking the body, walking over objects, and walking to music, but over time these strategies lose effectiveness [87].

BalanceTutor system allows for several unique postural control ability assessments as well as reactive response training in both standing and walking using controlled perturbation in simple task or multitask performance as follows:

6.1.3.1 Abnormal gait characteristics - Step forward initiation



Before perturbation




After perturbation

Patient position:	Standing comfortably with minimal external support
Platform's direction:	Anterior to posterior
Clinical effects:	Forward step initiation as part of gait training

6.1.3.2 Abnormal gait characteristics - stopping after starting




Before perturbation




After perturbation

Patient position:	Comfortable walking speed with minimal external support
Platform's direction:	Anterior to posterior
Comments:	This treatment is also recommended to be practiced while standing

6.1.3.3 Abnormal gait characteristics - shuffling steps




Before perturbation



After perturbation

Patient position:	Comfortable walking speed with minimal external support
Platform's direction:	Anterior to posterior
Comments:	Prolonged period of time at accelerated speed to Improve of stride length

6.1.3.4 Abnormal gait characteristics - Freezing of Gait




Before perturbation




After perturbation

Patient position:	Comfortable walking speed with minimal external support
Platform's direction:	Anterior to posterior
Comments:	Minimal period of time at highly accelerated perturbation in various walking speed

6.1.3.5 Falls Prevention - Equilibrium




Before perturbation




After perturbation

Patient position:	Standing comfortably with minimal external support
Platform's direction:	Medial/ Lateral
Comments:	Mild perturbation in order to a cause ankle and hip balance strategy while standing or walking

6.1.3.6 Falls prevention - Balance



Before perturbation




After perturbation

Patient position:	Comfortable walking speed with minimal external support
Platform's direction:	Medial/ Lateral
Comments:	Customized challenge forcing the initiation of a compensatory step while standing or walking

6.1.3.7 Increase walking speed




Before perturbation



After perturbation

Patient position:	Comfortable walking speed with minimal external support
Platform's direction:	Anterior to posterior
Comments:	Increasing walking speed- Challenging the patient to reach max ability

6.1.3.8 Abnormal gait characteristics - Neuromuscular coordination



Before perturbation



After perturbation

Patient position:	Comfortable walking speed with minimal external support
Platform's direction:	Random utilizing four directions
Comments:	Coordination of joint movements and muscle activation

6.1.4 Falls Prevention


Falls are a major problem among elderly population; they are the leading cause of injury above the age of 65 [88]. Of those who fall in the U.S., 20 to 30 % suffer moderate to severe injuries that reduce mobility and independence, and increase the risk of death [89]. The financial impact in 2000, adjusted for inflation, was \$30 billion and is expected to reach \$67.7 billion by 2020 [88]. Walking is the major activity in which large proportion of falls in older adults occurs [90]. Sixty percent of outdoor falls among older adults resulted from slips or trips [91]. Even among older adults capable of independent walking, there could be a substantial decline in balance performance, which does not become evident until a slip or a trip happens [92]. In fact, the inability to step rapidly in response to unexpected loss of balance ultimately determines whether a fall occurs [93, 94]. Thus, a better way to improve balance, improve stepping and reduce risk of falls may be to direct preventive efforts towards older adults who have not yet fallen. Until recently these balance recovery responses were considered hardwired postural reflexes that could not be influenced by training. However, [95–97] it was showed that older adults were able to adapt in a reactive manner after participation in a perturbation exercises that challenged the mechanisms responsible for dynamic stability (i.e., increase in base of support and counter-rotating segments around the center of mass). A number of studies have begun to examine the effect of perturbation training on balance of older adults. Shimada

et al. [98] found improvement in mobility and a trend to fall reduction after split treadmill training. This training method is a very unnatural condition, given that most people walk with the same velocity in each leg. Other studies have other issues. Pai et al. [99] showed a rapid decrease in loss of balance in response to multiple presentations of a slip perturbation after rising from sit to stand. Mansfield et al [100] found that older adults with a history of falls or instability reduced the frequency of multi-step reactions and foot collisions after perturbation training while standing or walking in place. The training methods perturb the balance of their participants from sit to stand or during standing or walking in place, which may not be as relevant in a natural setting as might be a perturbation while walking. Melzer and Oddsson [101] found improvement in voluntary stepping, and balance control, in an exercises program that incorporate mild external balance perturbation exercises applied by the instructors; and Halvarsson et al. [102] found that old fallers that suffered from fear of falling, decreased their fear of falling, and voluntary stepping times during dual-task performance and increased velocity of walking post perturbation training.. However, in these program the perturbations of posture were expected, and not random. Recently it was showed that unidirectional translational treadmill training (i.e., a laboratory induced trip) reduced falls [103, 104]. Bhatt et al. [105] found that inducing unannounced right-leg slips, participants significantly reduced fall and balance loss incidence. Pai, et al. [106] found that a single session of repeated-slip exposure reduced older adults' annual risk of falls from 34 to 15 % ($p < 0.05$) especially among those who had history of falls. The above protocol provided an anterior perturbation, causing a backwards "slip" initiated always by on the right foot. Participants might have learned and expected the right-leg slips perturbations. In a recent meta-analysis [107] that include 8 perturbation-based balance training studies ($n = 404$) participants reported fewer falls than those in the control groups. Motivated by the above perturbations training studies and trying to accommodate for some of the issues mentioned above (i.e., perturbation training while standing or walking in place; highly predictable repeated-right leg slip exposure; a very unnatural walking on split treadmill), we aimed to explore whether unexpected multidirectional perturbation training while walking on a treadmill [108] can reduce risks of falls in independent older adults. A perturbation exercise while walking provides a more realistic balance training that is sufficiently task-specific so that responses on this training regime will be more likely to be transferred to other measures of balance control and voluntary stepping measures. Exposure to a gait training program that includes unexpected perturbations exercises during walking will significantly improve voluntary stepping times as well as balance control in older adults, two factors that are associated with falls and injuries related with falls [109–114]. Perturbation exercises


that will challenge both balance control during walking as well as trigger a quick stepping responses to avoid fall during walking. This postural response following an external perturbation receives a higher priority than a voluntary action thus can be incorporated into centrally programmed voluntary movements [115]. This concept should be of importance for balance training and it further supports the notion that postural perturbations should be incorporated into balance training programs.

BalanceTutor system allows for several unique postural control ability assessments as well as reactive response training in both standing and walking using controlled perturbation in simple task or multitask performance as follows:

6.1.4.1 Forward automatic postural response



Before perturbation




After perturbation

Patient position: Standing comfortably


Platform's direction: Anterior to Posterior

Clinical effects: Fast and highly intensive activation of lower limb flexors muscles

6.1.4.2 Backward automatic postural response



Before perturbation




After perturbation

Patient position: Standing comfortably


Platform's direction: Posterior to Anterior

Clinical effects: Fast and highly intensive activation of Lower limb extensors muscles

6.1.4.3 Left side automatic postural response



Before perturbation




After perturbation

Patient position: Standing comfortably


Platform's direction: Medial to Lateral

Clinical effects: Fast and highly intensive activation of Left limb

6.1.4.4 Right side automatic postural response



Before perturbation




After perturbation

Patient position: Standing comfortably


Platform's direction: Lateral to Medial

Clinical effects: Fast and highly intensive activation of Right limb

6.1.4.5 Dynamic balance training



Before perturbation



After perturbation

Patient position: Walking or running

Platform's direction: Random Multi-Direction perturbation

Clinical effects: Faster compensatory step
Improvement of Recovery time

6.2 Orthopedic and Sport Rehabilitation

Rehabilitation continues to evolve with the increased emphasis on patient management and proprioceptive training. Proprioception can be defined as a specialized variation of the sensory modality of touch that encompasses the sensation of joint movement (kinesthesia) and joint position (joint position sense). Numerous investigators have observed that afferent feedback to the brain and spinal pathways is mediated by skin, articular, and muscle mechanoreceptors. Examining the effects of ligamentous injury, surgical intervention, and proprioceptive mediated activities in the rehabilitation program provides an understanding of the complexity of this system responsible for motor control. It appears that this neuromuscular feedback mechanism becomes interrupted with injury and abnormalities, and approaches restoration after surgical intervention and rehabilitation. Rehabilitation programs should be designed to include a proprioceptive component that addresses the following three levels of motor control: spinal reflexes, cognitive programming, and brainstem activity. Such a program is highly recommended to promote dynamic joint and functional stability. Thus far, current knowledge regarding the basic science and clinical application of proprioception has led the profession of sports medicine one step closer to its ultimate goal of restoring function. The proper management of athletic-related injuries and orthopedic lesions can be complex in the sports medicine settings. One of the most challenging aspects to the clinician is the understanding the role of proprioceptively mediated neuromuscular control after joint injury and its restoration through rehabilitation. Proprioception contributes to the motor programming for neuromuscular control required for precision movements and also contributes to muscle reflex, providing dynamic joint stability. The coupling effect of ligamentous trauma resulting in mechanical instability and proprioceptive deficits contributes to functional instability, which could ultimately lead to further micro trauma and occurrence of re-injury. Improvement of functional and sport-specific activities after musculoskeletal trauma and rehabilitation can be enhanced significantly if proprioception is addressed and introduced early in the treatment program. In addition to the mechanical restraint provided by articular structures, it has been observed that ligaments provide neurologic feedback that directly mediates reflex muscular stabilization around the joint. The inclusion of proprioception in the rehabilitation program should be based on the preceding findings and not on anecdotal evidence without an understanding of the neuromuscular mechanism. This understanding, coupled with a base of knowledge regarding the current research on proprioception, is necessary for sports medicine practitioners to optimize treatment programs for athletes. Numerous researchers have provided definitions regarding the terminology of joint sensation, or proprioception and kinesthesia [118,141].

Most contemporary authorities define proprioception as a specialized variation of the sensory modality of touch that encompasses the sensation of joint movement (kinesthesia) and joint position (joint position sense). The sensory receptors for proprioception that are found in the skin, muscles, and joints as well as in ligaments and tendons all provide input to the central nervous system (CNS) regarding tissue deformation [130]. Visual and vestibular centers also contribute afferent information to the CNS regarding body position and balance [148]. Trauma to tissues, that contain mechanoreceptors, may result in partial differentiation, which can lead to proprioceptive deficits. Susceptibility to re-injury, therefore, becomes a realistic possibility because of this decrease in proprioceptive feedback. However, studies have shown at least partial restoration of kinesthesia and joint position sense in surgically reconstructed shoulders and knees after rehabilitation [139,140]. Regaining neuromuscular control after injury or surgery is a necessary prerequisite for athletes wishing to return to competition. The neural input that is provided by the peripheral mechanoreceptors as well as the visual and vestibular receptors are all integrated by the CNS to generate a motor response. These responses generally fall under three levels of motor control: spinal reflexes, cognitive programming, and brainstem activity. In a situation where a joint is placed under mechanical loading, reflex muscular stabilization is stimulated through the spinal reflexes[132]. Cognitive programming that involves the highest level of CNS function (motor cortex, basal ganglia, and the cerebellum refers to voluntary movements that are repeated and stored as central commands. This awareness of body position and movement allows various skills to be performed without continuous reference to consciousness [148]. As defined earlier, proprioceptive feedback plays a major role in the conscious and unconscious awareness of a joint or limb in motion. The concept of proprioception is based on the fact that neural feedback to the CNS is mediated by cutaneous, muscle and joint mechanoreceptors. When examining the neural composition of joints, Hilton's law states that joints are innervated by articular branches of the nerves supplying the muscles that cross the joint[134]. In addition to proprioceptive mechanoreceptors, articular structures also include nociceptive free nerve endings. Activation of joint mechanoreceptors are triggered by the deformation and loading of the soft tissues that compose the joint. This neural stimulation travels to the CNS for integration via cortical and reflex pathways. These mechanoreceptors demonstrate adaptive properties depending on a particular stimulus [130]. Quick-adapting joint mechanoreceptors, such as the Pacinian corpuscles, decrease their discharge rate to extinction within milliseconds of the onset of a continuous stimulus. The Ruffini ending, Ruffini corpuscles, and the Golgi tendon-like organs that are referred to as the slow adapting mechanoreceptors, continue their discharge in

response to a continuous stimulus [130]. The properties of the quick-adapting mechanoreceptors lead to the notion that they mediate the sensation of joint motion because they are very sensitive to changes in position. Muscular mechanoreceptors and Ruffini ending joint receptors are slow adapting mechanoreceptors and are thought to mediate the sensation of joint position and changes in position because they are maximally stimulated at specific joint angles. One form of the slow-adapting receptors is the complex, fusiform muscle spindle receptors found within skeletal muscle. The muscle spindle receptor functions to measure muscle tension over a large range of extrafusal muscle length. It has been suggested that muscle and joint mechanoreceptors are complementary to each other in providing afferent input in regard to limb position [130].

This relationship between muscle and joint mechanoreceptors has been supported by the identification of the neural components necessary for the sensation of motion (rapidly adapting receptors, e.g., Pacinian corpuscles), joint position and acceleration (slow-adapting receptors, e.g., Ruffini endings and Ruffini corpuscles), and pain (free nerve endings) within ligamentous, cartilaginous, and muscular structures of the joints.

The spindle receptors found within the muscle are composed of a small bundle of modified muscle fibers called intrafusal fibers, to which the endings of several sensory nerves are attached. The extrafusal fibers that form the bulk of the muscle are responsible for generating force and are innervated by the alpha-motor neurons whereas the intrafusal fibers are innervated by the gamma-motor neurons. In the case of muscle contraction, alpha-gamma co-activation is believed to be the mechanism through which muscle length and tension are monitored. Activation of gamma-motor neurons allows the readjustment of spindle sensitivity in the case where extrafusal fibers are shortened. This allows the spindles to be functional at all times during a contraction. When a muscle is loaded beyond an anticipated level, intrafusal fiber shortening occurs to a greater degree than extrafusal shortening. Stretching of the spindles in the central region causes a burst of excitatory postsynaptic potentials from spindle afferents. These signals summate with the alpha-motor neurons from descending pathways, thereby increasing force production [142].

With respect to changes in muscle and tendon tension, the Golgi tendon organs function as a protective mechanism. The Golgi tendon organs are located within the tendons of the muscle and are

recruited when muscle contraction pulls on the tendon, which straightens the collagen bundles and distorts the receptor endings of the afferent neurons^[142].

This distortion increases the discharge rate of action potentials of these receptors that travel and synapse on spinal interneurons that project to motor neurons. Increased activity of the Golgi tendon organ afferents result in the inhibition of the motor neurons innervating the muscles that were stretched while exciting the motor nerves of the antagonistic muscles.

Studies have demonstrated the detrimental effect on reflex joint stabilization as a result of joint injury ^[145,146]. The contribution of musculotendinous receptors over joint receptors to the proprioceptive reflex remains a controversial issue in the literature.

Clinical application of proprioception research findings must be achieved at various levels of care for the management of orthopedic lesions. A thorough understanding of proprioception assessment techniques will aid the clinician and orthopedic surgeon to apply what is currently known concerning the non-traumatized joint to muscular, ligamentous, and cartilaginous injury. Establishing the effects of musculoskeletal trauma on joint position sensibility and neuromuscular control assists the need for critical decision-making regarding the appropriateness of various forms of treatment. From this point, it is known that surgical intervention plays an important role in the restoration of mechanical joint stability but its effects on proprioception pathways require further clarification. Finally, rehabilitation activities incorporating principles that stimulate the different levels of motor control to facilitate the return to function should be examined not only for its theoretical basis but also for its practical applicability.

The assessment of neuromuscular control includes the measurement of cortical, spinal reflex, and brainstem pathways. The evaluation of this complex neuromuscular system as different components allows a more detailed explanation of afferent control mechanisms. As defined previously, kinesthesia and joint position sense are components of proprioception. Functionally, kinesthesia is assessed by measuring threshold to detection of passive motion while joint position sense is assessed by measuring reproduction of passive positioning and reproduction of active positioning. When tested at a slow angular velocity (0.5 to 2 deg/sec), threshold to detection of passive motion as well as the reproduction of passive positioning is thought to selectively stimulate Ruffini or Golgi-type

mechanoreceptors. Because the test is performed passively, it is believed to maximally stimulate joint receptors, thereby relying on the cortical pathway in the neuromuscular control system. After ligament lesions, passive joint sensibility testing is often chosen to assess afferent activity because muscle activity is negated. Stimulation of both joint and muscle receptors is done by the reproduction of active positioning, which provides a more functional assessment of the afferent pathways. The evaluation of reflex capabilities is often assessed by measuring the latency of muscular activation to involuntary perturbation via Electromyography interpretation. The ability to quantify the sequencing of muscle firing can provide a valuable tool for the assessment of asynchronous neuromuscular activation patterns that may predispose an articulation to overuse trauma.

Functional assessment of the combined peripheral, vestibular, and visual contributions to neuromuscular control is best accomplished through the use of balance and postural sway measurements for the lower extremity. The availability of stabilometric methods and instrumentation can provide a relatively accurate index for these measures.

To reiterate this understanding of the proprioception mechanism, the clinician must apply the available knowledge to try to delineate the effects of orthopedic injury, surgical reconstruction, and rehabilitation on the various afferent pathways. Specifically, clinical research aimed at determining the effects of injury, surgery, and rehabilitation on joint position sensibility, neuromuscular control as well as balance and postural sway can provide a solid foundation for the development of a testing model for the knee, ankle, and shoulder that attempt to address these issues.

Numerous studies have been performed that examined the role of proprioception in the knee joint. It has been found that damage to articular structures, such as the ACL and meniscus, in addition to osteoarthritic changes disrupts articular structures containing mechanoreceptors. The disruption in the cortical pathway, therefore, results in an alteration in joint position sense and kinesthesia. Barrack et al. ~ and Skinner et al^[143].observed decreased kinesthesia with increasing age and ACL disruption. Joint position sensibility decrements have also been documented as a result of osteoarthritic changes in the knee^[117].

Deficits in the neuromuscular reflex pathway may have a detrimental effect on this motor control system's role as a protective mechanism in acute knee injury. The initiation of the reflex arc stimulated by mechanoreceptors and muscle spindle receptors occurs at a faster rate than signals induced by nociceptors (70 to 100 m/sec versus 1 m/sec) [130]. This suggests that proprioception may play a more significant role than pain impulses in preventing injury in the acute setting. However, the incidence of reinjury and the cause of chronic injuries may be attributed, to a greater extent, to proprioceptive deficits. These deficits may be induced by partial deafferentation as a result of initial knee injury and may also contribute to chronic joint disease through a decrease in joint afferents. This phenomenon has been observed by Beard et al [120]. in subjects with arthroscopically confirmed ACL deficiency. A significant deficit in reflex activation of the hamstring muscles after a 100 N anterior shear force in a single legged closed kinetic chain position was identified, as compared with the contralateral uninjured limb. Furthermore, Solomonow et al [145]. found that a direct stress applied to the ACL resulted in reflex hamstring activity, thereby contributing to the maintenance of joint integrity.

Although Barrack et al [116]. demonstrated a proprioceptive deficit after ACL disruption, it appears that kinesthetic awareness may be partially restored after ACL reconstruction. Kinesthesia has been reported to be restored after surgery as detected by the threshold to the detection of passive motion in the midrange of motion (45°). However, a longer threshold to the detection of passive motion was observed in the ACL reconstructed knee compared with the contralateral uninvolved knee when tested at 15° of flexion. Lephart et al [139]. found similar results in patients after either arthroscopically assisted patellar-tendon autograft or allograft ACL reconstruction. This evidence suggests that kinesthesia may have returned in the midrange of motion after ACL reconstruction and appears to be more sensitive in the near-terminal range of motion. The importance of incorporating a proprioceptive element in any comprehensive rehabilitation program is justified based on the results of these studies. Proprioceptive deficits may predispose an athlete to reinjury through decrements in the neuromuscular pathways resulting in the inhibition of complete rehabilitation.

Chronic ankle instability as a result of partial deafferentation of articular mechanoreceptors with joint injury was first postulated by Freeman et al. [10]. They observed that a decrease in the ability to maintain a one-legged stance occurred in the sprained ankle versus the contralateral uninjured ankle. The effect of unilateral ankle sprains on cortical pathway measures of proprioception have been

investigated by Garn and Newton, [127] who measured the ability of a subject to properly sense a passive movement or no movement state in the sagittal plane. Deficits in the ability to actively replicate passive ankle and foot positioning in this plane was reported by Glencross and Thornton [128] while testing the sprained ankle versus the contralateral uninjured ankle. Gross [131] recently reported that an increased probability of reinjury occurs as a result of a decrease in sensory input from joint receptors, leading to abnormal body positioning and diminished postural reflex responses. It was also found by Konradsen and Ravn [136] that chronic ankle instability resulted in a prolonged peroneal reaction time in response to a sudden inversion stress when compared with age-matched controls. Partial deafferentation resulting in diminished reflex joint stabilization may contribute to these findings.

The development of high technologic systems to assess the effects of musculoskeletal injury on balance has occurred in an attempt to quantify both static and dynamic components of proprioception. [133] The method of evaluation is based on the notion that damage to joint proprioceptors after injury to the lateral ligamentous complex of the ankle diminishes afferent feedback from the injured joint, thereby resulting in increases in postural sway [125].

To date, however, documented evidence exists concerning the alterations in postural sway after ankle injury using subjective evaluation (i.e., Romberg test). No increases in postural sway were observed by Tropp and Odenrick [147] when comparing a group of soccer players with previous ankle sprains to a control group of uninjured soccer players. Furthermore, no differences in postural sway were found between the involved and uninvolved ankles in a group of soccer players with a history of unilateral, recurrent ankle sprains. [147].

However, significant increases in postural sway were observed by Cornwall and Murrell [123] when comparing patients with acute ankle sprains with uninjured controls as long as 2 years after their injuries.

The effects of surgical reconstruction for functional ankle instability on proprioception pathways as measured through joint position sensibility or balance and postural sway assessments have not yet been thoroughly investigated. Empirical evidence exists suggesting that proprioceptive training techniques after acute and chronic ankle injuries are highly effective. In addition, ankle wrapping

and bracing have also been suggested to have a proprioceptive benefit. However, this notion remains untested and, therefore, unproven.

The objectives of proprioceptive rehabilitation are to retrain altered afferent pathways to enhance the sensation of joint movement. Proprioceptively mediated neuromuscular control of joints takes into account three distinct levels of motor activation within the CNS. Reflexes at the spinal level mediate movement patterns that are received from higher levels of the nervous system. This action provides for reflex joint stabilization during conditions of abnormal stress about the articulation and has significant implications for rehabilitation.^{2°} The use of exercises that facilitate dynamic joint stabilization may result in the improvement of this neuromuscular mechanism. The second level of motor control, located within the brainstem, receives input from joint mechanoreceptors, vestibular centers, and visual input from the eyes to maintain posture and balance of the body. Reactive neuromuscular activities that allow this pathway to process input from the aforementioned forms of afferent stimuli can be used to enhance brainstem function.

The highest level of CNS function (motor cortex, basal ganglia, and cerebellum) provides cognitive awareness of body position and movement in which motor commands are initiated for voluntary movements. Use of the cortical pathway allows movements that are repeated and stored as central commands to be performed without continuous reference to consciousness. Kinesthetic and proprioception training are such types of activity that can enhance this function.

Incorporating the three levels of motor control into activities to address proprioceptive deficiencies should be initiated early during the rehabilitation process [138]. Encouraging maximum afferent discharge to the respective CNS level must be the goal in stimulating joint and muscle receptors. [148]. To stimulate reflex joint stabilization, which emanates from the spinal cord, activities should focus on sudden alterations in joint positioning that necessitate reflex neuromuscular control. Enhancing motor function at the brainstem level can be achieved by performing balance and postural activities, both with and without visual input. Maximally stimulating the conversion of conscious to unconscious motor programming can be achieved by performing joint positioning activities, especially at joint end ranges. [148]. Simple tasks such as balance training and joint repositioning should begin early in the rehabilitation program and should become increasingly more difficult as the patient progresses. Regaining joint sense awareness to initiate muscular reflex stabilization to

prevent reinjure should be the primary objective once the final stage of rehabilitation is reached. Some contemporary authors believe that adaptations that occur during rehabilitation are related to (mediated by) feed-forward processing and are less a function of enhanced afferent pathways [137]. This theory suggests that fast movements are controlled by advance information known about the task, while concurrent proprioceptive feedback is relatively less important. Feedback is used primarily at the cortical level to determine the success or failure of that movement and to a lesser extent at the subcortical level for directing the movement. With repetition, the cerebral cortex can determine the most effective motor pattern for a given task, based on the proprioceptive information of previous attempts [137]. Biofeedback training appears to use the feed-forward learning process. However, there is still controversy regarding the contribution of afferent feedback in feed-forward processing. In order to achieve optimum rehabilitation the trainer has to accurately target and customize the practice to joint stability. Joint stability relies of neuromuscular coordination which in turn is dependent on proper proprioception function. This therefore singles out rehabilitation of the proprioception system as an important target during rehabilitation.

Exercises that focus on proprioceptive rehabilitation will improve postural control and balance ability. There are two main methods that target will result in an improvement in the target the In order to improve proprioception function balance practice and postural control training customized to the level of joint stabilization needs to be implemented.

Traditional balance practice works through anticipatory exercises. response ability so as to afford neuromuscular reeducation in all rehabilitation stages, covering both the inner and outer limits of ability from injury to return to play. In addition the practice has to be specific enough so that increments in difficulty levels do not lead to further injury.

Traditional rehabilitation relies on the use of proactive postural control practice. This means joint stabilization is performed as a result of a proactive movement in the body's center of mass. The trainer's ability to customize the practice is limited by both the limits and the specificity of the practice.

Perturbation is the introduction of an unexpected event that actively moves the body's centre of mass. Perturbation elicits a reactive or anticipatory postural control reflex like movement. Reactive

response training enforces the use of selective co-contraction patterns which keeps the joint within its physiological range of motion and avoids injury.

In order to better understand how anticipatory postural control practice more effectively targets proprioception a brief recap in the role of type I and type II mechanoreceptors is needed.

Type I mechanoreceptors have a low threshold and slow adaptation and are involved in proactive postural control. Type II mechanoreceptors have a low threshold and rapid adaptation and are more involved in responding to rapid unexpected joint destabilization elicited by perturbation and reactive postural control.


In order to avoid recurrent injury caused by a sudden perturbation to postural control it is important to train for this eventuality by practicing movements that safely require the input from type I and type II mechanoreceptors are necessary to avoid injury train for co-contraction It is also important Reactive response training by eliciting reactive response to perturbation both type I and Type II mechanoreceptors are forced to work and the proprioception system is optimized.

Proprioceptive system relies on wide spectrum and high resolution sensation using different types of specific Mechanoreceptors dealing with both expected and also following an ankle sprain, part of this specific mechanoreceptors mainly oriented rapid unexpected perturbation detectors at the earliest stage of the sprain event are damaged leading to recurrent ankle sprain.


At high ability level the use of reactive training will initiate the of type II mechanoreceptors which are necessary for reactive postural control. A higher level of reactive postural control ability and more efficient use of type II mechanoreceptors will decrease the risk of recurrent injury.

BalanceTutor system allows for several unique postural control ability assessments as well as proprioceptive rehabilitation using reactive response training in both standing and walking using controlled perturbation in simple task or multitask performance as follows:

6.2.1.1 ACL Rehabilitation




Before perturbation




After perturbation

Patient position:	Standing on injured leg with knee slightly bent
Platform's direction:	Anterior/ posterior
Clinical effects:	Fast and Highly Intensive activation of Knee muscles Use of ACL Proprioceptive and Kinesthesia abilities at maximum capacity Accurate and Fast Quadriceps and Hamstrings Coordination

6.2.1.2 Ankle extension Rehabilitation




Before perturbation



After perturbation

Patient position:	Standing comfortably
Platform's direction:	Posterior to Anterior
Clinical effects:	Fast and highly intensive activation of ankle extensors muscles Unique technique for rearfoot weight bearing High level coordination of ankle agonist and antagonist muscle activation

6.2.1.3 Ankle flexion rehabilitation



Before perturbation



After perturbation

Patient position:	Standing comfortably
Platform's direction:	Anterior to Posterior
Clinical effects:	<p>Fast and highly intensive activation of Gastrocnemius and Soleus muscles</p> <p>Unique technique for Forefoot weight bearing</p> <p>High level coordination of ankle agonist and antagonist muscle activation</p> <p>Unique facilitation of gait initiation</p>

6.2.1.4 Ankle instability rehabilitation



Before perturbation



After perturbation

Patient position:	Standing on injured leg with knee slightly bent
Platform's direction:	Multi-Direction Unexpected Perturbation
Clinical effects:	<p>High Level ankle neuromuscular coordination</p> <p>Static and Dynamic ankle joint stabilization</p> <p>High resolution of Proprioceptive and Kinesthesia activation</p>

6.2.1.5 Knee stabilization



Before perturbation



After perturbation

Patient position: Standing on injured leg with knee slightly bent

Platform's direction: Multi-Direction Unexpected Perturbation

Clinical effects: High Level ankle neuromuscular coordination
 Static and Dynamic ankle joint stabilization
 High resolution of Proprioceptive and Kinesthesia activation

6.2.1.6 Neuromuscular Coordination Training



Before perturbation



After perturbation

Patient position: Walking at customized speed

Platform's direction: Multi-Direction Unexpected Perturbation

Clinical effects: Fast Co-contraction of Agonist and Antagonist muscles of the limb in stance phase.
 Fast Co-ordination of Agonist and Antagonist muscles of the limb in swing phase

6.2.1.7 Groin strain rehabilitation




Before perturbation



After perturbation

Patient position:	Standing while both feet are together
Platform's direction:	Anterior to Posterior Perturbation while walking sideways
Clinical effects:	Controlled contraction and stretching of the hip adductors

6.2.1.8 Hip flexion rehabilitation




Before perturbation



After perturbation

Patient position:	Standing comfortably
Platform's direction:	Anterior to Posterior perturbation standing or walking
Clinical effects:	Unique technique for stretching of Iliopsoas in stance phase Fast contraction of Iliopsoas through automatic response at swing phase

6.2.1.9 Hip extension rehabilitation



Before perturbation




After perturbation

Patient position: Standing comfortably


Platform's direction: Posterior to Anterior Perturbation while standing and walking

Clinical effects: Fast contraction of Gluteus maximus

6.2.1.10 Hip abduction rehabilitation



Before perturbation



After perturbation

Patient position: Comfortable walking speed with minimal external support

Platform's direction: Medial/ Lateral Perturbation while standing or walking

Clinical effects: Fast contraction of Gluteus medius toward opposite direction of applied perturbation

6.2.1.11 Pure Reactive Response Training



Before perturbation



After perturbation

Patient position:	Walking or running
Platform's direction:	Random Multi-Direction perturbation
Clinical effects:	Improvement of Automatic Postural adjustment Faster compensatory step Improvement of Recovery time

6.2.1.12 Multi task training



Before perturbation




After perturbation

Patient position:	Standing, walking or running at customized speed
Platform's direction:	Multi-Direction perturbation
Clinical effects:	Neuromuscular coordination Fast Static and Dynamic multi-joints stabilization Optimal balance training

6.2.1.13 Weight bearing



Before perturbation




After perturbation

Patient position: Based on instructions of the game

Platform's direction: None, Shifting of Center of Pressure (COP)

Clinical effects: COP feedback training

6.2.1.14 Low back muscle strengthening



Before perturbation




After perturbation

Patient position: Standing comfortably

Platform's direction: Posterior to Anterior

Clinical effects: Fast and highly intensive activation of ankle extensors muscles
 Unique technique for Rear foot weight bearing
 High level coordination of ankle agonist and antagonist muscle activation

6.2.1.15 Abdominal muscle strengthening



Before perturbation



After perturbation

Patient position:	Standing comfortably
Platform's direction:	Anterior to Posterior
Clinical effects:	<p>Fast and highly intensive activation of Gastrocnemius and Soleus muscles</p> <p>Unique technique for Forefoot weight bearing</p> <p>High level coordination of ankle agonist and antagonist muscle activation</p> <p>Unique facilitation of gait initiation</p>

7 References

1. Holloway RG, Tuttle D, Baird T, Skelton WK. The safety of hospital stroke care. *Neurology*. 2007;68(8):550–55.
2. Davenport RJ, Dennis MS, Wellwood I, Warlow CP. Complications after acute stroke. *Stroke*. 1996;27(3):415–20.
3. Truelsen T, Piechowski-Józwiak B, Bonita R, Mathers C, Bogousslavsky J, Boysen G. Stroke incidence and prevalence in Europe: a review of available data. *Eur J Neurol*. 2006;13(6):581–98.
4. Carter KN, Anderson CS, Hackett ML, Barber PA, Bonita R; Auckland Regional Community Stroke Study Group. Improved survival after stroke: Is admission to hospital the major explanation? Trend analyses of the Auckland Regional Community Stroke Studies. *Cerebrovasc Dis*. 2007; 23(2–3):162–68.
5. Ramnemark A, Nilsson M, Borssen B, Gustafson Y. Stroke, a major and increasing risk factor for femoral neck fracture. *Stroke*. 2000;31(7):1572–77.
6. Ramnemark A, Nyberg L, Borssen B, Olsson T, Gustafson Y. Fractures after stroke. *Osteoporos Int*. 1998; 8(1):92–5
7. O’Loughlin JL, Robitaille Y, Boivin JF, Suissa S. Incidence of and risk factors for falls and injurious falls among the community-dwelling elderly. *Am J Epidemiol*. 1993;137(3):342–54.
8. Tinetti ME, Speechley M, Ginter SF. Risk factors for falls among elderly persons living in the community. *N Engl J Med*. 1988;319(26):1701–7.
9. Campbell AJ, Borrie MJ, Spears GF, Jackson SL, Brown JS, Fitzgerald JL. Circumstances and consequences of falls experienced by a community population 70 years and over during a prospective study. *Age Ageing*. 1990;19(2):136–41.
10. Rubenstein LZ. Falls in older people: epidemiology, risk factors and strategies for prevention. *Age Ageing*. 2006; 35 Suppl 2:ii37–ii41.
11. Tutuarima JA, Van der Meulen JH, De Haan RJ, Van Straten A, Limburg M. Risk factors for falls of hospitalized stroke patients. *Stroke*. 1997;28(2):297–301.
12. Smith J, Forster A, Young J. Use of the “STRATIFY” falls risk assessment in patients recovering from acute stroke. *Age Ageing*. 2006;35(2):138–43.
13. Mayo NE, Korner-Bitensky N, Kaizer F. Relationship between response time and falls among stroke patients undergoing physical rehabilitation. *Int J Rehabil Res*. 1990; 13(1):47–55.
14. Nyberg L, Gustafson Y. Fall prediction index for patients in stroke rehabilitation. *Stroke*. 1997;28(4):716–21.
15. Nyberg L, Gustafson Y. Patient falls in stroke rehabilitation. A challenge to rehabilitation strategies. *Stroke*. 1995; 26(5):838–42.
16. Nyberg L, Gustafson Y. Using the Downton index to predict those prone to falls in stroke rehabilitation. *Stroke*. 1996;27(10):1821–24.
17. Forster A, Young J. Incidence and consequences of falls due to stroke: a systematic inquiry. *BMJ*. 1995;311(6997): 83–86.
18. Sze KH, Wong E, Leung HY, Woo J. Falls among Chinese stroke patients during rehabilitation. *Arch Phys Med Rehabil*. 2001;82(9):1219–25.

19. Teasell R, McRae M, Foley N, Bhardwaj A. The incidence and consequences of falls in stroke patients during inpatient rehabilitation: factors associated with high risk. *Arch Phys Med Rehabil.* 2002;83(3):329–33.
20. Suzuki T, Sonoda S, Misawa K, Saitoh E, Shimizu Y, Kotake T. Incidence and consequence of falls in inpatient rehabilitation of stroke patients. *Exp Aging Res.* 2005; 31(4):457–69.
21. Aizen E, Shugaev I, Lenger R. Risk factors and characteristics of falls during inpatient rehabilitation of elderly patients. *Arch Gerontol Geriatr.* 2007;44(1):1–12.
22. Watanabe Y. Fear of falling among stroke survivors after discharge from inpatient rehabilitation. *Int J Rehabil Res.* 2005;28(2):149–52.
23. McLean DE. Medical complications experienced by a cohort of stroke survivors during inpatient, tertiary-level stroke rehabilitation. *Arch Phys Med Rehabil.* 2004; 85(3): 466–69.
24. Roth EJ, Lovell L, Harvey RL, Heinemann AW, Semik P, Diaz S. Incidence of and risk factors for medical complications during stroke rehabilitation. *Stroke.* 2001;32(2): 523–29.
25. Jørgensen L, Engstad T, Jacobsen BK. Higher incidence of falls in long-term stroke survivors than in population controls: depressive symptoms predict falls after stroke. *Stroke.* 2002;33(2):542–47.
26. Yates JS, Lai SM, Duncan PW, Studenski S. Falls in community-dwelling stroke survivors: an accumulated impairments model. *J Rehabil Res Dev.* 2002;39(3):385–94.
27. Hyndman D, Ashburn A. People with stroke living in the community: attention deficits, balance, ADL ability and falls. *Disabil Rehabil.* 2003;25(15):817–22.
28. Hyndman D, Ashburn A. Stops walking when talking as a predictor of falls in people with stroke living in the community. *J Neurol Neurosurg Psychiatry.* 2004;75(7):994–97.
29. Soyuer F, Ozturk A. The effect of spasticity, sense and walking aids in falls of people after chronic stroke. *Disabil Rehabil.* 2007;29(9):679–87.
30. Harris JE, Eng JJ, Marigold DS, Tokuno CD, Louis CL. Relationship of balance and mobility to fall incidence in people with chronic stroke. *Phys Ther.* 2005;85(2):150–58.
31. Mackintosh SF, Hill KD, Dodd KJ, Goldie PA, Culham EG. Balance score and a history of falls in hospital predict recurrent falls in the 6 months following stroke rehabilitation. *Arch Phys Med Rehabil.* 2006;87(12):1583–89.
32. Mackintosh SF, Hill K, Dodd KJ, Goldie P, Culham E. Falls and injury prevention should be part of every stroke rehabilitation plan. *Clin Rehabil.* 2005;19(4):441–51.
33. Belgen B, Beninato M, Sullivan PE, Narielwalla K. The association of balance capacity and falls self-efficacy with history of falling in community-dwelling people with chronic stroke. *Arch Phys Med Rehabil.* 2006;87(4):554–61.
34. Hyndman D, Ashburn A, Stack E. Fall events among people with stroke living in the community: circumstances of falls and characteristics of fallers. *Arch Phys Med Rehabil.* 2002;83(2):165–70.
35. Andersson AG, Kamwendo K, Seiger A, Appelros P. How to identify potential fallers in a stroke unit: validity indexes of 4 test methods. *J Rehabil Med.* 2006;38(3): 186–91.

36. Lamb SE, Ferrucci L, Volapto S, Fried LP, Guralnik JM; Women's Health and Aging Study. Risk factors for falling in home-dwelling older women with stroke: the Women's Health and Aging Study. *Stroke*. 2003;34(2):494–501.
37. Stapleton T, Ashburn A, Stack E. A pilot study of attention deficits, balance control and falls in the subacute stage following stroke. *Clin Rehabil*. 2001;15(4):437–44.
38. Tutuarima JA, De Haan RJ, Limburg M. Number of nursing staff and falls: a case-control study on falls by stroke patients in acute-care settings. *J Adv Nurs*. 1993;18(7): 1101–5.
39. Zdobysz JA, Boradia P, Ennis J, Miller J. The relationship between functional independence scores on admission and patient falls after stroke. *Top Stroke Rehabil*. 2005; 12(2):65–71.
40. Berg WP, Alessio HM, Mills EM, Tong C. Circumstances and consequences of falls in independent community dwelling older adults. *Age Ageing*. 1997;26(4):261–68.
41. Michael KM, Allen JK, Macko RF. Reduced ambulatory activity after stroke: the role of balance, gait, and cardiovascular fitness. *Arch Phys Med Rehabil*. 2005;86(8): 1552–56.
42. Bohannon RW. Number of pedometer-assessed steps taken per day by adults: a descriptive meta-analysis. *Phys Ther*. 2007;87(12):1642–50.
43. Ugur C, Gucuyener D, Uzuner N, Ozkan S, Ozdemir G. Characteristics of falling in patients with stroke. *J Neurol Neurosurg Psychiatry*. 2000;69(5):649–51.
44. Hyndman D, Ashburn A, Yardley L, Stack E. Interference between balance, gait and cognitive task performance among people with stroke living in the community. *Disabil Rehabil*. 2006;28(13–14):849–56.
45. Bloem BR, Steijns JA, Smits-Engelsman BC. An update on falls. *Curr Opin Neurol*. 2003;16(1):15–26.
46. Rapport LJ, Webster JS, Flemming KL, Lindberg JW, Godlewski MC, Brees JE, Abadee LS. Predictors of falls among right-hemisphere stroke patients in the rehabilitation setting. *Arch Phys Med Rehabil*. 1993;74(6):621–26.
47. Webster JS, Roades LA, Morrill B, Rapport LJ, Abadee PS, Sowa MV, Dutra R, Godlewski MC. Rightward orienting bias, wheelchair maneuvering, and fall risk. *Arch Phys Med Rehabil*. 1995;76(10):924–28.
48. Muro MJ, De Pedro-Cuesta J, Almazan J, Holmqvist LW. Stroke patients in south Madrid: Function and motor recovery, resource utilization, and family support. *Stroke*. 2000;31(6):1352–59.
49. Dennis MS, Lo KM, McDowall M, West T. Fractures after stroke: frequency, types, and associations. *Stroke*. 2002;33(3):728–34.
50. Wei TS, Hu CH, Wang SH, Hwang KL. Fall characteristics, functional mobility and bone mineral density as risk factors of hip fracture in the community-dwelling ambulatory elderly. *Osteoporos Int*. 2001;12(12):1050–55.
51. *Epidemiology of MS*. New York (NY): National Multiple Sclerosis Society; 2008 [cited 2008 Nov 21].
52. Kaplin AI, Williams M. How common are the “common” neurologic disorders? *Neurology*. 2007;69(4):410–11.
53. Bourdette DN, Prochazka AV, Mitchell W, Licari P, Burks J. Health care costs of veterans with multiple sclerosis: Implications for the rehabilitation of MS. VA Multiple Sclerosis Rehabilitation Study Group. *Arch Phys Med Rehabil*. 1993;74(1):26–31.
54. Cattaneo D, De Nuzzo C, Fascia T, Macalli M, Pisoni I, Cardini R. Risk of falls in subjects with multiple sclerosis. *Arch Phys Med Rehabil*. 2002;83(6):864–67.

55. Finlayson ML, Peterson EW, Cho CC. Risk factors for falling among people aged 45 to 90 years with multiple sclerosis. *Arch Phys Med Rehabil.* 2006;87(9):1274–79.
56. Nilsagård Y, Lundholm C, Denison E, Gunnarsson LG . Predicting accidental falls in people with multiple sclerosis A
57. longitudinal study. *Clin Rehabil.* 2009;23(3):259–69.
58. Chen LH, Warner M, Fingerhut L, Makuc D. Injury episodes and circumstances: National Health Interview Survey, 1997–2007. *Vital Health Stat* 10. 2009;(241):1–55.
59. Anderson RN, Miniño AM, Fingerhut LA, Warner M, Heinen MA. Deaths: Injuries, 2001. *Natl Vital Stat Rep.* 2004;52(21):1–86.
60. Hornbrook MC, Stevens VJ, Wingfield DJ, Hollis JF, Greenlick MR, Ory MG . Preventing falls among community dwelling older persons: Results from a randomized trial. *Gerontologist.* 1994;34(1):16–23.
61. Hausdorff JM, Rios DA, Edelberg HK. Gait variability and fall risk in community-living older adults: A 1-year prospective study. *Arch Phys Med Rehabil.* 2001;82(8):1050–56.
62. Gillespie LD, Robertson MC, Gillespie WJ, Lamb SE, Gates S, Cumming RG , Rowe BH. Interventions for preventing falls in older people living in the community. *Cochrane Database Syst Rev.* 2009;(2):CD007146.
63. Gillespie LD, Gillespie WJ, Robertson MC, Lamb SE, Cumming RG , Rowe BH. Interventions for preventing falls in elderly people. *Cochrane Database Syst Rev.* 2003;(4)
64. Guideline for the prevention of falls in older persons. American Geriatrics Society, British Geriatrics Society, and American Academy of Orthopaedic Surgeons Panel on Falls Prevention. *J Am Geriatr Soc.* 2001;49(5):664–72.
65. Injury prevention & control: Data & statistics (WISQARS) [Internet]. Atlanta (GA): Centers for Disease Control and Prevention; 2005
66. Hu G , Baker SP. Recent increases in fatal and non-fatal injury among people aged 65 years and over in the USA. *Inj Prev.* 2010;16(1):26–30.
67. Stevens JA, Corso PS, Finkelstein EA, Miller TR. The costs of fatal and non-fatal falls among older adults. *Inj Prev.* 2006;12(5):290–95.
68. Englander F, Hodson TJ, Terregrossa RA. Economic dimensions of slip and fall injuries. *J Forensic Sci.* 1996;41(5): 733–46.
69. Peterson EW, Cho CC, Von Koch L, Finlayson ML. Injurious falls among middle aged and older adults with multiple sclerosis. *Arch Phys Med Rehabil.* 2008;89(6):1031–37.
70. Brønnum-Hansen H, Hansen T, Koch-Henriksen N, Stenager E. Fatal accidents among Danes with multiple sclerosis. *Mult Scler.* 2006;12(3):329–32.
71. Peterson EW, Cho CC, Finlayson ML. Fear of falling and associated activity curtailment among middle aged and older adults with multiple sclerosis. *Mult Scler.* 2007;13(9): 1168–75.
72. Nilsagård Y, Denison E, Gunnarsson LG , Boström K. Factors perceived as being related to accidental falls by persons with multiple sclerosis. *Disabil Rehabil.* 2009;31(16): 1301–10.
73. Gray P, Hildebrand K. Fall risk factors in Parkinson’s disease. *J Neurosci Nurs.* 2000;32(4):222–28.
74. Morris ME. Movement disorders in people with Parkinson disease: A model for physical therapy. *Phys Ther.* 2000; 80(6):578–97.

75. PADRECC Home [homepage on the Internet]. Philadelphia (PA): U.S. Department of Veterans Affairs; c2001–07 [updated 2006 Nov 8; cited 2006 Aug 17]. Revolutionizing Parkinson's disease care for our veterans; [1 screen]. Available
76. Giladi N, McMahon D, Przedborski S, Flaster E, Guillory S, Kostic V, Fahn S. Motor blocks in Parkinson's disease. *Neurology*. 1992;42(2):333–39.
77. Giladi N, McDermott MP, Fahn S, Przedborski S, Jankovic J, Stern M, Tanner C; Parkinson Study Group. Freezing of gait in PD: Prospective assessment in the DATATOP cohort. *Neurology*. 2001;56(12):1712–21.
78. Lamberti P, Armenise S, Castaldo V, De Mari M, Iliceto G, Tronci P, Serlenga L. Freezing gait in Parkinson's disease. *Eur Neurol*. 1997;38(4):297–301.
79. Lewis GN, Byblow WD, Walt SE. Stride length regulation in Parkinson's disease: The use of extrinsic, visual cues. *Brain*. 2000;123(Pt 10):2077–90.
80. Weghorst S. Augmented reality and Parkinson's disease. *Commun ACM*. 1997;40(8):47–48.
81. Dietz MA, Goetz CG, Stebbins GT. Evaluation of a modified inverted walking stick as a treatment for Parkinsonian freezing episodes. *Mov Disord*. 1990;5(3):243–47.
82. Morris ME, Iansak R, Matyas TA, Summers JJ. Stride length regulation in Parkinson's disease. Normalization strategies and underlying mechanisms. *Brain*. 1996;119(Pt 2):551–68.
83. Fukunaga H, Kasai T, Yoshidome H. Clinical findings, status of care, comprehensive quality of life, daily life therapy and treatment at home in patients with Parkinson's disease. *Eur Neurol*. 1997;38 Suppl 2:64–69.
84. Schrag A, Jahanshahi M, Quinn NP. What contributes to depression in Parkinson's disease? *Psychol Med*. 2001;31(1): 65–73.
85. Karlsen KH, Tandberg E, Arslan D, Larsen JP. Health related quality of life in Parkinson's disease: A prospective longitudinal study. *J Neurol Neurosurg Psychiatry*. 2000; 69(5):584–89.
86. Defebvre L, Blatt JL, Blond S, Bourriez JL, Gieu JD, Destee A. Effect of thalamic stimulation on gait in Parkinson disease. *Arch Neurol*. 1996;53(9):898–903.
87. Stern GM, Lander CM, Lees AJ. Akinetic freezing and trick movements in Parkinson's disease. *J Neural Transm Suppl*. 1980;(16):137–41.
88. Centers for Disease Control and Prevention, National Center for Injury Prevention and Control. Web-based Injury Statistics Query and Reporting System (WISQARS) [online]. Accessed August 15, 2013.
89. Stevens JA. Falls among older adults—risk factors and prevention strategies. *J Saf Res*. 2005;36(4):409–11.
90. Robinovitch SN, Feldman F, Yang Y, et al. Video capture of the circumstances of falls in elderly people residing in long-term care: an observational study. *Lancet*. 2013;381(9860):47–54.
91. Luukinen H, Herala M, Koski K, et al. Fracture risk associated with a fall according to type of fall among the elderly. *Osteoporosis Int*. 2000;11(7):631–4.
92. Woollacott MH, Tang PF. Balance control during walking in the older adult: research and its implications. *Phys Ther*. 1997;77(6):646–60.
93. Maki BE, McIlroy WE. The role of limb movements in maintaining upright stance: the “change-in-support” strategy. *Phys Ther*. 1997;77(5):488–507.
94. Maki BE, McIlroy WE. Control of rapid limb movements for balance recovery: age-related changes and implications for fall prevention. *Age Ageing*. 2006; 35 Suppl 2:ii12–18.

95. Bierbaum S, Peper A, Karamanidis K, Arampatzis A. Adaptive feedback potential in dynamic stability during disturbed walking in the elderly. *J Biomech.* 2011;44(10):1921–6.
96. Bierbaum S, Peper A, Arampatzis A. Exercise of mechanisms of dynamic stability improves the stability state after an unexpected gait perturbation in elderly. *Age (Dordr).* 2013;35(5):1905–15. 10.
97. Arampatzis A, Peper A, Bierbaum S. Exercise of mechanisms for dynamic stability control increases stability performance in the elderly. *J Biomech.* 2011;44(1):52–8.
98. Shimada H, Obuchi S, Kamide N, et al. Relationship with dynamic balance function during standing and walking. *Am J Phys Med Rehabil.* 2003;82(7): 511–6.
99. Pai Y, Bhatt T, Wang E, et al. Inoculation against falls: rapid adaptation by young and older adults to slips during daily activities. *Arch Phys Med Rehabil.* 2010;91(3):452–9.
100. Mansfield A, Peters AL, Liu BA, et al. Effect of a perturbation-based balance training program on compensatory stepping and grasping reactions in older adults: a randomized controlled trial. *Phys Ther.* 2010;90(4):476–91.
101. Melzer I, Oddsson LI. Improving balance control and self-reported lower extremity function in community-dwelling older adults: a randomized control trial. *Clin Rehabil.* 2013;27(3):195–206.
102. Halvarsson A, Oddsson L, Olsson E, et al. Effects of new, individually adjusted, progressive balance group training for elderly people with fear of falling and tend to fall: a randomized controlled trial. *Clin Rehabil.* 2011; 25(11):1021–31.
103. Grabiner MD, Bareither ML, Gatts S, et al. Task-specific training reduces trip-related fall risk in women. *Med Sci Sports Exerc.* 2012;44(12):2410–4.
104. Rosenblatt NJ, Marone J, Grabiner MD. Preventing trip-related falls by community-dwelling adults: a prospective study. *J Am Geriatr Soc.* 2013;61(9): 1629–31.
105. Bhatt T, Yang F, Pai Y. Learning to resist gait-slip falls: long-term retention in community-dwelling older adults. *Arch Phys Med Rehabil.* 2012;93(4):557–64.
106. Pai YC, Bhatt T, Yang F, et al. Perturbation training can reduce community dwelling older adults' annual fall risk: a randomized controlled trial. *J Gerontol A Biol Sci Med Sci.* 2014;69(12):1586–94.
107. Shapiro A, Melzer I. Short report balance perturbation system to improve balance compensatory responses during walking in old persons. *J Neuroeng Rehabil.* 2010;7:32.
108. Mansfield A, Wong JS, Bryce J, Knorr S, Patterson KK. Does perturbation-based balance training prevent falls? Systematic review and meta-analysis of preliminary randomized controlled trials. *Phys Ther.* 2015;95(5):700–9.
109. Melzer I, Kurz I, Shahar D, et al. Predicting injury from falls in older adults: comparison of voluntary step reaction times in injured and noninjured fallers—a prospective study. *J Am Geriatr Soc.* 2009;57(4):743–5.
110. Melzer I, Kurz I, Shahar D, et al. Do voluntary step reactions in dual task conditions have an added value over single task for fall prediction? A prospective study. *Aging Clin Exp Res.* 2010;22(5-6):360–6.
111. Melzer I, Kurz I, Shahar D, et al. Application of the voluntary step execution test to identify elderly fallers. *Age Ageing.* 2007;36(5):532–7.
112. Melzer I, Benjuya N, Kaplanski J. Postural stability in the elderly: a comparison between fallers and non-fallers. *Age Ageing.* 2004;33(6):602–7.

113. Melzer I, Kurz I, Oddsson LI. A retrospective analysis of balance control parameters in elderly fallers and non-fallers. *Clin Biomech (Bristol, Avon)*. 2010;25(10):984–8.
114. Kurz I, Oddsson L, Melzer I. Characteristics of balance control in older persons who fall with injury—a prospective study. *J Electromyogr Kinesiol*. 2013;23(4):814–9.
115. Layne CS, Abraham LD. Interactions between automatic postural adjustments and anticipatory postural patterns accompanying voluntary movement. *Int J Neurosci*. 1991;61(3-4):241–54.
116. Collins JJ, De Luca CJ. Open-loop and closed-loop control of posture: a random-walk analysis of center-of-pressure trajectories. *Exp Brain Res*. 1993;95(2):308–18.
117. Barrack RL, Skinner HB, Buckley SL. Proprioception in the anterior cruciate deficient knee. *Am J Sports Med* 17 1-6, 1989
118. Barrett DS, Cobb AG, Bentley G. Joint proprioception in normal, osteoarthritic and replaced knees. *J Bone Joint Surg* 73B. 53-56, 1991
119. Bastian HC. The "muscular sense", its nature and cortical localization. *Brain* 10 1-137, 1988
120. Baxendale RH, Ferrell WR, Wood L. Responses of quadriceps motor units to mechanical stimulation of knee joint receptors in the decerebrate goat. *Brain Res* 453 150-156, 1988
121. Beard DJ, Kyberd PJ, O'Connor JJ, et al. Reflex hamstring contraction latency in anterior cruciate ligament deficiency. *J Orthop Res* 12- 219- 228, 1994
122. Borsa PA, Lephart SM, Kocher MS, et al: Functional assessment and rehabilitation of shoulder proprioception for glenohumeral instability. *J Sport Rehab* 3 84-104, 1994
123. Boyd IA. The histological structure of the receptors in the knee joint of the cat correlated with their physiological response. *J Physiol* 124 476-488, 1954
124. Cornwall MW, Murrell P. Postural sway following inversion sprain of the ankle. *J Am Podiatr Med Assoc* 81 243-247, 1991
125. Dunn TG, Gillig SE, Ponsor SE, et al. The learning process in biofeedback : Is it feed-forward or feedback? *Biofeedback Self-Regul* 11 143- 156, 1986
126. Freeman MAR, Dean M, Hanham I: The etiology and prevention of functional instability of the foot. *J Bone Joint Surg* 47B 669-677, 1965
127. Freeman MAR, Wyke B. The innervation of the knee joint. An anatomical and histological study in the cat. *J Anat* 101 505-532, 1964
128. Garn SN, Newton RA. Kinesthetic awareness in subjects with multiple ankle sprains. *Phys Ther* 68 1667-1671, 1988
129. Glencross D, Thornton E. Position sense following joint injury. *J Sports Med Phys Fitness* 21 23-27, 1981
130. Gloumann R, Jobe FW, Tibone JE, et al. Dynamic electromyographic analysis of the throwing shoulder with glenohumeral instability. *J Bone Joint Surg* 70A 220-226, 1988
131. Gngg P. Peripheral neural mechanisms in proprioception. *J Sport Rehab*
132. Gross MT. Effects of recurrent lateral ankle sprains on active and passive judgements of joint position. *Phys Ther* 67 1505-1509, 1987
133. Guanche C, Knatt T, Solomonow M, et al: The synergistic action of the capsule and the shoulder muscles. *Am J Sport Med* 23 301-306, 1995

134. Guskiewicz KM, Perrin DH: Research and clinical applications of assessing balance J Sport Rehabil 5 45-63, 1996
135. Hilton J On the Influence of Mechanical and Physiological Rest in the Treatment of Accidents and Surgical Diseases and the Diagnostic Value of Pain A Course of Lectures London, Bell & Daldy, 1863
136. Kennedy JC, Alexander IJ, Hayes KC Nerve supply of the human knee and its functional importance. Am J Sports Med 10 329-335, 1982
137. Konradsen L, Ravn JB. Ankle instability caused by prolonged peroneal reaction time Acta Orthop Scand 61 388-390, 1990
138. Leisman G Cybernetic model of psychophysiological pathways: II Consciousness of tension and kinesthesia J Manipulative Physiol Ther 12 174-191, 1989
139. Lephart SM, Henry TJ The physiological basis for open and closed kinetic chain rehabilitation for the upper extremity. J Sport Rehab 5 71-87, 1996
140. Lephart SM, Kocher MS, Fu FH, et al Proprioception following ACL reconstruction. J Sport Rehab 1 186-196, 1992
141. Lephart SM, Warner JP, Borsa PA, et al Proprioception of the shoulder in normal, unstable and post-surgical individuals J Shoulder Elbow Surg 3 371-380, 1994
142. McCloskey DI Kinesthetic sensibility Physiol Rev 58 763-820, 1978
143. Moffett DF, Moffett SB, Schauf CL Human Physiology. Foundations and Frontiers. St Louis, Mosby-Year Book, Inc., 1993
144. Skinner HB, Barrack RL, Cook SD. Age-related decline in proprioception. Clin Orthop 184 208-211, 1984
145. Smith RL, Brunolli J Shoulder kinesthesia after shoulder dislocation Phys Ther 69 106-112, 1989
146. Solomonow M, Baratta R, Zhou BH, et al. The synergistic action of the anterior cruciate ligament and thigh muscles in maintaining joint stability. Am J Sports Med 15. 207-213, 1987
147. Tibone JE, Antich TJ, Fanton GS, et al. Functional analysis of anterior cruciate ligament instability. Am J Sports Med 14: 276-284, 1986
148. Tropp H, Odenrick P Postural control in single-limb stance J Orthop Res 6 833-839, 1988
149. Tyldesling B, Greve Ji Muscles, Nerves and Movement Kinesiology in Daily Living Boston, Blackwell Scientific Publications, 1989, pp 268-284
150. Willis WD, Grossman RG Medical Neurobiology Neuroanatomical and Neurophysiological Principles Basic to Clinical Neuroscience Third edition St. Louis, CV Mosby Co , 1981, pp 123-128

8 Version History

Date	Change description	Written by	Ver.
5 Oct 2016	Initial revision	Dr. Avraham Cohen.	161005